

Patient profile

Enterprise One received 100 completed referral questionnaires in this period (all services, n = 473)

Demographic information	Your service	All services
Sex (female)	72.0%	67.9%
Average age (years)	12.8	12.7
Aboriginal and/or Torres Strait Islander	4.3%	7.1%
Living in highest disadvantage area	16.8%	16.8%

Clinical characteristics (carer reported)	Your service	All services
Patients using medication daily	65.2%	60.3%
Patients using opioid medication daily (excluding codeine)	4.4%	3.1%
Outside healthy weight range	48.0%	41.5%
Patients experiencing pain > 12-months	63.0%	61.8%
School days missed in last 2 weeks (Avg)	4.2	3.9
Carer work impairment due to child's pain (Average %)	43.6%	45.4%

Clinical characteristics (patient reported)	Your service	All services	
Pain severity	5.3	5.2 📕	
	Mild / Moderate / Severe		
PedsQL Total - Psychosocial	46.1 50.4	47.1 50.9	
- Physical - Sleep	37.9 29.8	39.7 33.8	
% patients "at risk" of impaired health related quality of life	93.2%	91.4%	
Functional disability	27.1	26.2	
No or m	ninimal / Modera	ate / Severe	
BAPQ Pain-related worry	16.0	15.9	

Episode information

	Your service	All services
Episodes of care started	89	463
Waiting time* (median days)	36.5	35.5
No. of treatment pathways started	69	359
Pathway delivery mode (%) - In clinic care	92.6%	92.4%
- Remote care	4.4%	5.9%
Treatment pathway type (%)		
- Group program	15.9%	13.6%
- Individual	81.2%	82.7%
Episodes of care ended	72	330
Ave length of episode (lapsed days)	324.3	320.5

* Days from referral received to episode start

Collection of patient outcomes

A detailed summary of outcome collection rates at episode end and at post-episode follow-up can be found in the **appendices of your report**.

Indicator and benchmark summary


	Your service		All services	
Benchmark or indicator	Patient outcomes included (n)	Positive outcomes* (%)	Patient outcomes included (n)	Positive outcomes* (%)
Time from referral to episode start (benchmark) Percentage of episodes started within 8 weeks of referral being received	88	64.8%	458	63.1%
Average pain (indicator) Percentage of patients with moderate or worse 'average pain' at referral who make a clinically significant improvement at episode end	13	38.5%	58	50.0%
Paediatric quality of life (indicator) Percentage of patients with impaired health related quality of life at referral who make a clinically significant improvement at episode end	15	80.0%	75	80.0%
Functional disability (indicator) Percentage of patients with at least moderate functional disability at referral who make a clinically significant improvement at episode end	15	46.7%	65	53.8%


* Positive outcomes describes the percentage of patients making a clinically significant improvement, or reducing oMEDD by 50% or

having their episode start within 3 months/8 weeks where relevant


- is shown for the benchmarks and indicators where less than 10 outcomes are included

Time from referral to episode start (benchmark)






Paediatric quality of life (indicator)




Patients with impaired health related quality of life at referral who make a clinically significant improvement at episode end (%)

Average pain (indicator)



Patients with at least moderate average pain at referral who make clinically significant improvement at episode end (%)

Functional disability (indicator)



Patients with at least moderate functional disability at referral who make clinically significant improvement at episode end (%)